

University of Dundee

Phosphatase-mediated bioprecipitation of lead by soil fungi

Liang, Xinjin; Kierans, Martin; Ceci, Andrea; Hillier, Stephen; Gadd, Geoffrey Michael

Published in:
Environmental Microbiology

DOI:
[10.1111/1462-2920.13003](https://doi.org/10.1111/1462-2920.13003)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Liang, X., Kierans, M., Ceci, A., Hillier, S., & Gadd, G. M. (2016). Phosphatase-mediated bioprecipitation of lead by soil fungi. *Environmental Microbiology*, 18(1), 219-231. <https://doi.org/10.1111/1462-2920.13003>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Table 1. Tolerance indices (TI), pH values, P_i release and Pb-removal from MCD liquid media after growth of *Aspergillus niger* and *Paecilomyces javanicus*.

	TI (%)		pH				P _i (mM)				Pb concentration in supernatant (mM)		Pb removed by fungal biomass (μmol ml ⁻¹)		Pb removed from solution (%)	
	G2P+Pb	PyA+Pb	G2P	PyA	G2P+Pb	PyA+Pb	G2P	PyA	G2P+Pb	PyA+Pb	G2P+Pb	PyA+Pb	G2P+Pb	PyA+Pb	G2P+Pb	PyA+Pb
MCD+																
Abiotic control	NA	NA	6.0 ± 0.0	6.0 ± 0.0	6.0 ± 0.0	6.0 ± 0.0	ND	ND	ND	ND	4.98 ± 0.02	4.94 ± 0.04	NA	NA	NA	NA
<i>A. niger</i> 10d	67.7 ± 0.6 %	56.3 ± 0.2 %	3.3 ± 0.2	3.9 ± 0.0	4.6 ± 0.1	3.7 ± 0.1	14.1 ± 0.6	5.9 ± 0.5	9.0 ± 0.5	9.9 ± 0.3	1.35 ± 0.18	1.38 ± 0.02	3.52 ± 0.13	3.58 ± 0.02	70.7	72.5
<i>A. niger</i> 20d	58.8 ± 0.7 %	78.0 ± 1.8 %	3.6 ± 0.0	3.5 ± 0.1	3.3 ± 0.3	3.4 ± 0.0	18.0 ± 1.7	6.1 ± 0.4	19.3 ± 0.8	15.4 ± 0.4	0.68 ± 0.02	0.44 ± 0.02	4.31 ± 0.13	4.56 ± 0.03	86.5	92.3
<i>A. niger</i> 30d	54.7 ± 0.6 %	108.1 ± 1.1 %	3.7 ± 0.1	3.5 ± 0.2	3.1 ± 0.0	3.3 ± 0.4	24.2 ± 0.8	7.6 ± 1.4	24.7 ± 0.8	26.0 ± 0.8	0.007 ± 0.02	0.005 ± 0.002	4.94 ± 0.04	4.92 ± 0.01	99.8	99.5
<i>P. javanicus</i> 10d	97.4 ± 0.2 %	59.1 ± 0.3 %	7.6 ± 0.1	6.1 ± 0.2	6.3 ± 0.1	5.9 ± 0.0	7.2 ± 0.5	6.1 ± 0.7	15.7 ± 1.7	13.8 ± 1.2	2.22 ± 0.03	1.47 ± 0.09	2.68 ± 0.15	3.52 ± 0.09	53.8	71.3
<i>P. javanicus</i> 20d	89.6 ± 0.3 %	144.8 ± 2.4 %	6.5 ± 0.3	6.3 ± 0.1	6.5 ± 0.2	6.2 ± 0.2	11.5 ± 0.8	8.4 ± 1.2	21.6 ± 1.9	15.5 ± 1.5	0.14 ± 0.28	1.31 ± 0.04	4.84 ± 0.18	3.66 ± 0.03	97.2	74.1
<i>P. javanicus</i> 30d	122.8 ± 0.5 %	177.6 ± 0.1 %	6.9 ± 0.1	6.6 ± 0.3	6.8 ± 0.4	6.6 ± 0.5	15.0 ± 0.4	9.5 ± 0.5	25.7 ± 0.7	25.9 ± 1.0	0.028 ± 0.03	0.03 ± 0.02	4.97 ± 0.01	4.91 ± 0.02	99.8	99.4

Tolerance indices (TI) were derived from the biomass dry weight ratios of lead-exposed mycelium to control mycelium harvested at 10, 20 and 30 d (Sayer *et al.*, 1995; Wei *et al.*, 2013). The pH values, and P_i and Pb concentrations are those obtained from analysis of culture supernatants after removal of biomass. Pb removed by the biomass is expressed as $\mu\text{mol ml}^{-1}$ culture medium for ease of comparison with supernatant Pb concentrations: values can be multiplied by 100 to give total μmol Pb removed. *A. niger* and *P. javanicus* were grown in MCD liquid media without Pb, or amended with 5 mM $\text{Pb}(\text{NO}_3)_2$, and containing 30 mM $\text{C}_3\text{H}_7\text{Na}_2\text{O}_6\text{P}\cdot\text{xH}_2\text{O}$ (G2P) or 5 mM $\text{C}_6\text{H}_{18}\text{O}_{24}\text{P}_6\cdot\text{xNa}^+\cdot\text{yH}_2\text{O}$ (PyA) as sole source of P. Organisms were grown at 25°C in the dark on an orbital shaking incubator at 125 rpm. Values shown are means of at least three measurements with typical relative standard deviations of about 5%. NA = not applicable; ND = not detectable.